

EFFECT OF SPACING AND DETOPPING ON DHAINCHA SEED PRODUCTION

A. J. Shinde¹, D. D. Mankar², S. S. Patil³, N. R. Mairan⁴ and M. N. Wairagade⁵

ABSTRACT

The study entitled “Effect of spacing and detopping on dhaincha seed production” was conducted during *kharif* 2019 in split plot design with three spacings as main plot treatments (S₁- 60 cm × 10 cm, S₂- 90 cm × 10 cm, S₃- 120 cm × 10 cm) and four detopping as sub plot treatments (D₀- Control- no topping, D₁- detopping at 30 DAS, D₂- detopping at 45 DAS and D₃- detopping at 60 DAS) with three replications.

The spacing of 60 cm × 10 cm (S₁) recorded significantly higher dry matter, pod weight and seed yield (1045.7 kg ha⁻¹), stalk yield (8190.2 kg ha⁻¹), GMR (73198 Rs ha⁻¹), NMR (56709 Rs ha⁻¹) and B: C ratio (4.43) over the other spacing treatments viz., 90 cm × 10 cm (S₂) and 120 cm × 10 cm (S₃).

Detopping at 30 DAS (D₁) recorded significantly higher plant height, number of branches plant⁻¹, dry matter, number of pods plant⁻¹, pod weight plant⁻¹, seed yield plant⁻¹, seed yield (923.4 kg ha⁻¹), stalk yield plant⁻¹ and stalk yield (7101.5 kg ha⁻¹), GMR (64639 Rs ha⁻¹), NMR (48460 Rs ha⁻¹) and B: C ratio (3.98) over the other detopping treatments viz., Control (D₀) and detopping at 60 DAS (D₃). However, detopping at 45 DAS (D₂) was found to be at par with detopping at 30 DAS (D₁)

(Keywords: Dhaincha, spacing, detopping, yield, economics)

INTRODUCTION

Dhaincha (*Sesbania aculeata* Pers.) is a leguminous crop, large genus of herbs and shrubs. It is used as animal feed and for the green manuring. Dhaincha crop makes good growth within short period about 2-4 months (for green manuring), having straight stem up to 6 m in height which is very soft in nature, the leaflets of crop are very tender and produces maximum amount of organic matter. In this respect dhaincha is a potential green manure crop which can produce up to 22.5 tonnes ha⁻¹ of biomass, the highest among all green manure crops and has nutrient content 3.3%N, 0.7% P, and 1.3%K (Chandra, 2005). It is annual crop with lower C: N ratio. Khan (1952) reported that 12-15% of yield of rice was increased by green manure with dhaincha crop. Roots of dhaincha crop produce large number of bacterial nodules through which fixes large amount of atmospheric nitrogen in the soil. Besides green manuring, it is also used as cover crop. It protects soil from water and wind erosion.

Seed production of dhaincha will not only improves the soil health status by adding nutrients (nitrogen) and organic matter but it will also generate the extra income to farmers. The lack of availability of adequate quality seeds at appropriate time at reasonable price for small holdings and

marginal farmers becomes a major constraint in dhaincha cultivation. Quality seed production of dhaincha is given meager importance in spite of huge demand from farmers.

It is known fact that optimum spacing provides sufficient light to all the plants which increases the rate of photosynthesis and consequently dry matter production. In case of *Sesbania aculeata* proper spacing provides sufficient interception of light and satisfactory absorption of nutrients and water from the soil due to the proper development of root system and results in higher crop yield.

Detopping is the act of mowing or cutting the aerial part of a crop. Detopping of terminal buds is carried out two months after sowing to encourage more branching along with improve seed yield. Removal of apical dominance promotes the development of lateral buds thereby resulting in increased branches plant⁻¹ (Pathania *etal.*, 2000). Hence, the experiment was conducted with the objectives to find out the optimum spacing, optimum time period of detopping for better branching with their economics.

MATERIALS AND METHODS

A field experiment was carried out at Agronomy Section Farm, College of Agriculture, Nagpur during *kharif* season of 2019-20 to study the “Effect of spacing and

- 1 & 5. P.G. Students, Agronomy Section, College of Agriculture, Nagpur (M.S.)
2. Assoc. Professor of Agronomy, ARS, Tharsa, dist. Nagpur (M.S.)
3. P.G. Student, SSAC Section, College of Agriculture, Nagpur (M.S.)
4. Sr. Res. Asstt., Agronomy Section, College of Agriculture, Nagpur (M.S.)

detopping on dhaincha seed production.” The soil of the experimental plot was clayey in texture, low in available nitrogen (205.18 kg ha⁻¹), medium in available phosphorus (17.16 kg ha⁻¹) and very high in available potash (420.45 kg ha⁻¹) with pH 7.7. The experiment was conducted during *khari* 2019 in split plot design with three spacings as main plot treatments (S₁- 60 cm × 10 cm, S₂- 90 cm × 10 cm, S₃- 120 cm × 10 cm) and four detopping as sub plot treatments (D₀- Control- no topping, D₁- detopping at 30 DAS, D₂- detopping at 45 DAS and D₃- detopping at 60 DAS) with three replications. Local variety of dhaincha was sown at spacing as per the treatments. A uniform dose of 25: 25: 00 kg NPK ha⁻¹ was applied to all the treatments. Similarly all the detopping practices were imposed as per treatments. No accountable incidence of any disease and insect pest was observed during the crop life period. The data was recorded as per the standard procedure. Observations on plant height plant⁻¹, number of branches plant⁻¹ and dry matter were recorded at harvest and similarly observations on number of pods plant⁻¹, pod weight plant⁻¹, seed yield plant⁻¹ and ha⁻¹, stalk yield plant⁻¹ and ha⁻¹ and test weight were also recorded. GMR, NMR and B: C ratio were also calculated.

RESULTS AND DISCUSSION

Effect of spacing levels

Data in Table 1 revealed that, the spacing of 60 cm × 10 cm (S₁) recorded higher plant height and number of branches plant⁻¹ over the spacing of 90 cm × 10 cm (S₂) and 120 cm × 10 cm (S₃), while the spacing of 120 cm × 10 cm (S₃) recorded significantly higher dry matter accumulation plant⁻¹ (g) over the spacing of 60 cm × 10 cm (S₁), however, it was at par with spacing of 90 cm × 10 cm (S₂) at harvest. The spacing of 120 cm × 10 cm also recorded the highest value of yield attributing characters viz., number of pods plant⁻¹ (g), pod weight plant⁻¹ (kg), seed yield plant⁻¹ (g) and ha⁻¹ (kg) and stalk yield plant⁻¹ (g) and ha⁻¹ ((kg) at harvest, over the spacing of 60 × 10 cm (S₁) and found to be at par with spacing of 90 cm × 10 cm (S₂). The better performance of plants at wider row spacing may be attributed to least inter plant competition and greater availability of growth resources i. e. light, moisture and nutrients. The results are in close agreements with the findings of Parlawar *et al.* (2001). They reported that the growth attributed viz., leaves plant⁻¹, number of branches plant⁻¹, number of nodules plant⁻¹ were more under wider row spacing of 60 cm. However, the plant height was more with closer row spacing of 30 cm. Tripathi *et al.* (2013) reported that the individual plant performance with respect to plant height, basal diameter, dry matter accumulation, number of branches plant⁻¹ and yield attributes was found maximum under the spacing of 45 cm × 20 cm, whereas total biomass and seed yield unit⁻¹ area was obtained highest with the spacing of 30 cm × 20 cm.

The spacing of 60 cm × 10 cm (S₁) recorded significantly higher seed yield (1045.7 kg ha⁻¹), stalk yield

(8190.2 kg ha⁻¹), GMR (73198 Rs ha⁻¹), NMR (56709 Rs ha⁻¹) and B: C ratio (4.43) over the other spacing treatments viz., 90 cm × 10 cm (S₂) and 120 cm × 10 cm (S₃). Wider spacing recorded higher value of vegetative and reproductive growth parameters of individual plant over narrow spacing. However, in terms of seed yield ha⁻¹, stalk yield ha⁻¹, GMR ha⁻¹ and NMR ha⁻¹ narrow spacing found superior over wider spacing, it might be due to the fact that plant stand ha⁻¹ was higher over wider spacing and this higher number of plants in area gave higher yield. The results are in confirmation with the findings of Venkanna *et al.* (2013). They revealed that higher seed yield ha⁻¹, GMR, NMR and B: C ratio were recorded at row spacing of 30 cm closely followed by that from 45 cm, while the lowest values recorded at wider row spacing of 60 cm.

Effect of detopping levels

Among the different detopping practices the plant height at harvest was significantly influenced by the control treatment i.e. non topping (D₀) and recorded the maximum and significantly higher plant height and was at par with by detopping at 60 DAS (D₃). Taller plant was registered with no topping practice, it might be due to the fact that plants were not topped and as such plant grows to their original height without reduction. While, the growth characters i.e. number of branches plant⁻¹, dry matter accumulation plant⁻¹ (g) and yield attributing characters i.e. number of pods plant⁻¹, pod weight plant⁻¹ (g), seed yield plant⁻¹ and ha⁻¹, stalk yield plant⁻¹ and ha⁻¹ and test weight (g) recorded significantly higher values by detopping at 30 DAS (D₁) over detopping at 60 DAS (D₃) and control (D₀). However, it was at par with detopping at 45 DAS (D₂). The increment in yield attributes under the influence of topping might be attributed to better growth characteristics like dry matter accumulation plant⁻¹ coupled with profuse branching. Results obtained are in accordance with those recorded by Mandal *et al.* (2017). They reported that pinching had significant effect on plant height, number of branches plant⁻¹, number of pods plant⁻¹, pod yield plant⁻¹ (g) and seed yield plant⁻¹ and ha⁻¹. Kolhe *et al.* (2019) revealed that nipping at 60 DAS was found significantly superior over nipping at 90 DAS in respect to plant height, number of branches plant⁻¹ and dry matter accumulation plant⁻¹.

Among the different detopping treatments, detopping at 30 DAS (D₁) recorded significantly higher seed yield (923.4 kg ha⁻¹), stalk yield (7101.5 kg ha⁻¹), GMR (64639 Rs ha⁻¹), NMR (48460 Rs ha⁻¹) and B: C ratio (3.98) over the other detopping treatments viz., Control (D₀) and detopping at 60 DAS (D₃). However, detopping at 45 DAS (D₂) was found to be at par with detopping at 30 DAS (D₁). These results are in agreement with Siddagangamma *et al.* (2018). They reported that GMR, NMR and B: C ratio were obtained significantly superior at terminal bud nipping at 35 and 50 DAS over no terminal bud nipping. Kolhe *et al.* (2020) revealed that GMR, NMR and B: C ratio were obtained when nipping was done at 60 DAS over the nipping at 90 DAS and rest of the treatments.

Interaction effect

Interaction effect between different spacing and detopping levels on growth, yield attributing characters, yield and economics was found to be non significant.

From the results it is inferred that spacing of 60 cm × 10 cm and detopping at 30 to 45 DAS found to be the best for highest seed yield (kg ha⁻¹) in dhaincha. Spacing of 60 cm × 10 cm recorded highest gross monetary returns (Rs ha⁻¹), net monetary returns (Rs ha⁻¹) and B: C ratio in dhaincha. Detopping at 30 DAS recorded highest gross monetary returns (Rs ha⁻¹), net monetary returns (Rs ha⁻¹) and B: C ratio in dhaincha and detopping at 45 DAS was at par.

REFERENCES

- Chandra, K. 2005. Organic manure, Regional centre of organic farming, Bangalore pp.13.
- Khan, A.A. 1952. Dhaincha is an ideal green manure crop Indian Farming, **2** (3) :8
- Kolhe, R. R., N. D. Parlawar, D. J. Jiotode, V. S. Khawale, T. A. Chavhan and R. I. Samrutwar, 2019. Effect of nipping and growth retardant on growth, yield and uptake of nutrient of pigeonpea. J. Soil and Crops, **29** (2): 376- 380.
- Kolhe, R. R., N. D. Parlawar, D. J. Jiotode, V. S. Khawale, T. A. Chavhan and R. I. Samrutwar, 2020. Influence of nipping and growth retardant on yield and yield parameters and economics of pigeonpea. J. Soil and Crops, **30**(1): 99- 102.
- Mandal, S., A. Dutta and P.K.Bhattacharya , 2017.Effect of pinching and foliar nutrition on growth and yield of dhaincha (*Sesbania rostrata* Bremek & Oberm). Int. J. Curr. Res.**9**(10):59676-59681.
- Parlawar, N.D., D.G. Giri, R.M. Adapawar and P.V. Yadgirwar, 2001. Influence of seed rate, row spacing and phosphate level on seed yield and economics of dhaincha (*Sesbania aculeata* L.). PKV Res. J.**25**(2):68-72.
- Pathania, N. S., O. P. Sehgal and Y. C. Gupta, 2000.Pinching for flower regulation in sim carnation.J.Ornamental Hort. **3**: 114- 117.
- Siddagangamma, K. R., A. A. Choudhary, S. N. Potkile, D. G. Sonune and M. P. Punse, 2018. Effect of terminal bud nipping and salicylic acid spray on growth and yield of sesame. J. Soil and Crops, **28**(1): 216- 220
- Tripathi, M.K., Babita Chaudhary, S.R. Singh and H.R. Bhandari, 2013. Growth and yield of sunhemp (*Crotalaria juncea* L.) as influenced by spacing and topping practices. African J. Agric.Res.**8**(28): 3744-3749.
- Venkanna, B., B. Joseph, K.B. Suneetha Devi and A. Sivasankar,2013. Influence of row spacing and phosphorous levels on seed production of dhaincha (*Sesbania aculeata*). J.Res. ANGRAU. **41**(3): 124-126.

Rec. on 25.10.2020 & Acc. on 10.11.2020